

Scattering of non-relativistic charged-particles by electromagnetic radiation

M. Apostol

Department of Theoretical Physics, Institute of Atomic Physics,

Magurele-Bucharest MG-6, POBox MG-35, Romania

email: apoma@theory.nipne.ro

Abstract

The cross-section is computed for non-relativistic charged particles (like electrons, ions) scattered by electromagnetic radiation confined to a finite region (like the focal region of optical laser beams); for convenience, a potential scattering is included. The cross-section exhibits maxima at scattering angles given by the energy and momentum conservation in multi-photon absorption or emission processes. The scattering process addressed in this paper is distinct from the process dealt with in previous studies, where the scattering is immersed in the radiation field.

PACS: 34.50.Rk; 34.80.Qb; 03.65.Nk; 61.05.J-; 79.20.Ws

Key words: scattering, non-relativistic charged particles, electromagnetic radiation

The potential scattering of charged particles in the presence of electromagnetic radiation has enjoyed much interest in the past,[1]-[9] and is still enjoying.[10] In the context of lasers development these studies revealed the potential scattering assisted by multiple-photon exchange, as shown by the well-known Kroll-Watson cross-section.[1, 2] In the formulation of this problem the scattering is immersed in the radiation field, *i.e.* the asymptotic incoming and outgoing particle states and the scattering potential are included in the region containing the radiation. The duration of the laser pulse is much longer than the scattering time. The starting point of these approaches is the standard non-relativistic hamiltonian where the particle momentum $\mathbf{p} = m\mathbf{v} + e\mathbf{A}/c$ includes the electromagnetic contribution $e\mathbf{A}/c$ beside the purely mechanical contribution $m\mathbf{v}$ (the notations are the usual ones, *i.e.* m and e denote the particle mass and, respectively, charge, \mathbf{v} is the particle velocity, \mathbf{A} is the vector potential of the radiation field and c denotes the speed of light in vacuum). The Kroll-Watson cross-section corresponds to radiation-assisted potential scattering, *i.e.* it shows how the potential cross-section is modified by radiation; it becomes zero when the potential is removed.

With the advent of high-intensity lasers and strongly focused laser beams,[11]-[18] it appears the possibility of scattering charged particles by the radiation field confined to the focal region of the beam (the radiation is vanishing smoothly outside the focal region). In this case the asymptotic scattering states are radiation free; they are eigenstate of the quantum-mechanical momentum corresponding to the purely mechanical momentum, without including the electromagnetic contribution. This is the scattering problem addressed in this paper. It resembles to some extent the electron diffraction from standing light waves, where the scattering proceeds by spontaneous emission of Compton photons (Kapitsa-Dirac effect).[19] We envisage charged particles like electrons or ions, with non-relativistic energies, scattered off electromagnetic radiation confined to the focal

region (in vacuum) of an optical laser beam. Usually, the dimension of the focal region is of the order a few tens of radiation wavelengths and the radiation field has a reasonably high intensity, such that the non-relativistic character of the particle motion be preserved. For convenience, we include in the focal region a static potential. We assume first a laser pulse much longer than the radiation wavelength.

We consider a non-relativistic particle with mass m and charge e scattered by an electromagnetic radiation field with the vector potential $\mathbf{A} = \mathbf{A}_0 \cos(\omega t - \mathbf{k}\mathbf{r})$ (linear polarization), where \mathbf{A}_0 is the amplitude and ω and \mathbf{k} are the radiation frequency and, respectively, wavevector; t and \mathbf{r} denote time and, respectively, position. Since the phase velocity of the non-relativistic charge is much smaller than the speed of light c in vacuum ($\omega = ck$), we may neglect the spatial phase $\mathbf{k}\mathbf{r}$ in comparison with the temporal phase ωt ; consequently, the vector potential may be approximated by $\mathbf{A} \simeq \mathbf{A}_0 \cos \omega t$. The hamiltonian of the particle in the radiation field is the dipole hamiltonian

$$H = \frac{1}{2m}p^2 - e\mathbf{r}\mathbf{E}(t) + V(\mathbf{r}) \quad , \quad (1)$$

where $\mathbf{E}(t) = \mathbf{E}_0 \sin \omega t$, $\mathbf{E}_0 = \omega \mathbf{A}_0 / c$, is the electric field and $V(\mathbf{r})$ is a potential which does not depend on time. The non-relativistic character of the motion is preserved in the radiation field provided $eA_0/mc^2 \ll 1$. Making use of the notations $H_0 = p^2/2m$ and $U = -e\mathbf{r}\mathbf{E}$ the wavefunction ψ_i of the incident particle satisfies the Schrodinger equation

$$i\hbar \frac{\partial \psi_i}{\partial t} = (H_0 + U + V)\psi_i \quad (2)$$

with the initial condition (incoming state)

$$\psi_i^0 = \frac{1}{\sqrt{v}} e^{-\frac{i}{\hbar} E_i t + \frac{i}{\hbar} \mathbf{p}_i \mathbf{r}} \quad , \quad (3)$$

where \mathbf{p}_i is the initial momentum, E_i is the initial energy and v denotes the volume; the solution of equation (2) is given by

$$\psi_i = \psi_i^0 - \frac{i}{\hbar} e^{-\frac{i}{\hbar} H_0 t} \int_{-\infty}^t dt' e^{\frac{i}{\hbar} H_0 t'} (U + V) \psi_i \quad . \quad (4)$$

The wavefunction of the final scattering state (outgoing state) is

$$\psi_f = \frac{1}{\sqrt{v}} e^{-\frac{i}{\hbar} E_f t + \frac{i}{\hbar} \mathbf{p}_f \mathbf{r}} \quad , \quad (5)$$

where \mathbf{p}_f is the final momentum and E_f is the final energy. The transition amplitude (S -matrix) reads

$$a_{fi} = -\frac{i}{\hbar} \int_{-\infty}^{+\infty} dt (\psi_f, (U + V)\psi_i) \quad . \quad (6)$$

In equation (2) we insert successively the Goeppert-Mayer transform[20]

$$\begin{aligned} \psi_i &= e^{iS_1} \phi_i \quad , \quad S_i = -\frac{1}{\hbar} \int_{-\infty}^t dt' U(t') = -\frac{e}{\hbar c} \mathbf{r} \mathbf{A} \quad , \\ i\hbar \frac{\partial \phi_i}{\partial t} &= \left[\frac{1}{2m} (\mathbf{p} - \frac{e}{c} \mathbf{A})^2 + V \right] \phi_i \end{aligned} \quad (7)$$

and the Kramers-Henneberger transform[21]-[23]

$$\begin{aligned}\phi_i &= e^{iS_2} \chi_i, \quad S_2 = \frac{e}{\hbar mc} \int_{-\infty}^t dt' \mathbf{p} \mathbf{A} - \frac{e^2}{2\hbar mc^2} \int_{-\infty}^t dt' A^2 = \\ &= \frac{e}{\hbar mc \omega} \mathbf{p} \mathbf{A}_0 \sin \omega t - \frac{e^2}{8\hbar mc^2 \omega} A_0^2 (\sin 2\omega t + 2\omega t), \\ i\hbar \frac{\partial \chi_i}{\partial t} &= \left(H_0 + \tilde{V} \right) \chi_i, \quad \tilde{V}(\mathbf{r}) = V(\mathbf{r} - e\mathbf{E}/m\omega^2).\end{aligned}\tag{8}$$

We recognize in equations (7) the standard non-relativistic hamiltonian of the particle in the radiation field. As regards the potential V we limit ourselves to the Born approximation; consequently, the wavefunction χ_i can be written as

$$\chi_i = \psi_i^0 - \frac{i}{\hbar} e^{-\frac{i}{\hbar} H_0 t} \int_{-\infty}^t dt' e^{\frac{i}{\hbar} H_0 t'} \tilde{V} \psi_i^0 \tag{9}$$

and the transition amplitude becomes

$$\begin{aligned}a_{fi} &= -\frac{i}{\hbar} \int_{-\infty}^{+\infty} dt (\psi_f, (U + V) e^{iS_1} e^{iS_2} \psi_i^0) + \\ &+ \left(-\frac{i}{\hbar}\right)^2 \int_{-\infty}^{+\infty} dt (\psi_f, U e^{iS_1} e^{iS_2} e^{-\frac{i}{\hbar} H_0 t} \int_{-\infty}^t dt' e^{\frac{i}{\hbar} H_0 t'} \tilde{V} \psi_i^0) .\end{aligned}\tag{10}$$

The U -term in equation (10) corresponds to the scattering by the radiation field; we can see in equation (10) that the scattering by the potential V is dressed with radiation (terms $\sim V e^{iS_1} e^{iS_2}$, $\sim \tilde{V}$); in addition, there appears an interference term, which includes the product $U\tilde{V}$.

We may make certain simplifications in equation (10). It is easy to see that the phase S_1 (equations (7)) is of the order $(r/\lambda)(eA_0/\hbar\omega) \gg eA_0/\hbar\omega$, where λ is the radiation wavelength; while the phase S_2 (equations (8)) is of the order $(p/mc, eA_0/mc^2)(eA_0/\hbar\omega) \ll eA_0/\hbar\omega$; consequently, we may neglect the phase S_2 in equation (10). In addition, the position given by the argument of \tilde{V} (equations (8)) is of the order $r - \lambda_c(eA_0/\hbar\omega)$, where $\lambda_c = \hbar/mc$ is the Compton wavelength of the particle; it takes the potential \tilde{V} far away from its short range in very short times, especially for (reasonably) high-intensity radiation; similarly, the Coulomb potential is rapidly reduced to an appreciable extent by the radiation, such that we may neglect the potential \tilde{V} in equation (10).

Making use of these simplifications the transition amplitude given by equation (10) becomes

$$\begin{aligned}a_{fi} &\simeq -\frac{i}{\hbar} \int_{-\infty}^{+\infty} dt (\psi_f, U e^{iS_1} \psi_i^0) - \frac{i}{\hbar} \int_{-\infty}^{+\infty} dt (\psi_f, V e^{iS_1} \psi_i^0) = \\ &= \int_{-\infty}^{+\infty} dt (\psi_f, \left(\frac{\partial}{\partial t} e^{iS_1}\right) \psi_i^0) - \frac{i}{\hbar} \int_{-\infty}^{+\infty} dt (\psi_f, V e^{iS_1} \psi_i^0) ,\end{aligned}\tag{11}$$

or

$$\begin{aligned}a_{fi} &= \frac{1}{v} \int_{-\infty}^{+\infty} dt d\mathbf{r} e^{\frac{i}{\hbar}(E_f - E_i)t} \frac{\partial}{\partial t} e^{-\frac{ie}{\hbar\omega} \mathbf{r} \mathbf{E}_0 \cos \omega t} e^{\frac{i}{\hbar} \mathbf{p} \mathbf{r}} - \\ &- \frac{i}{\hbar v} \int_{-\infty}^{+\infty} dt d\mathbf{r} e^{\frac{i}{\hbar}(E_f - E_i)t} V e^{-\frac{ie}{\hbar\omega} \mathbf{r} \mathbf{E}_0 \cos \omega t} e^{\frac{i}{\hbar} \mathbf{p} \mathbf{r}} ,\end{aligned}\tag{12}$$

where $\mathbf{p} = \mathbf{p}_i - \mathbf{p}_f$ is the momentum transfer. In equation (12) we use the decomposition

$$e^{-\frac{ie}{\hbar\omega} \mathbf{r} \mathbf{E}_0 \cos \omega t} = \sum_{n=-\infty}^{+\infty} (-i)^n J_n(er\mathbf{E}_0/\hbar\omega) e^{-in\omega t} , \tag{13}$$

where J_n are Bessel functions (n being any integer), and get

$$a_{fi} = -\frac{2\pi i}{v} \sum_{n=-\infty}^{+\infty} (-i)^n \delta(E_f - E_i - n\hbar\omega) \int d\mathbf{r} [n\hbar\omega + V(\mathbf{r})] J_n(er\mathbf{E}_0/\hbar\omega) e^{\frac{i}{\hbar} \mathbf{p} \mathbf{r}} ; \tag{14}$$

we can see the occurrence of multiple-photon scattering processes with energy conservation $E_f = E_i + n\hbar\omega$. The transition probability per unit time is given by

$$w_{fi} = \frac{2\pi}{\hbar v^2} \sum_{n=-\infty}^{+\infty} \delta(E_f - E_i - n\hbar\omega) \left| \int d\mathbf{r} [n\hbar\omega + V(\mathbf{r})] J_n(e\mathbf{r}\mathbf{E}_0/\hbar\omega) e^{\frac{i}{\hbar}\mathbf{p}\mathbf{r}} \right|^2 ; \quad (15)$$

we multiply w_{fi} by the density of final states $vp_f^2 dp_f d\Omega / (2\pi\hbar)^3$, where $d\Omega$ is the element of solid angle, divide by the current density v_i/v , where v_i is the initial velocity, and integrate over the final momentum p_f in order to get the differential cross-section for the n -process

$$d\sigma_n = \frac{p_{fn}}{p_i} \left| \frac{m}{2\pi\hbar^2} \int d\mathbf{r} [n\hbar\omega + V(\mathbf{r})] J_n(e\mathbf{r}\mathbf{E}_0/\hbar\omega) e^{\frac{i}{\hbar}\mathbf{p}\mathbf{r}} \right|^2 d\Omega ; \quad (16)$$

the momentum p_{fn} is given by the energy conservation $p_{fn}^2/2m = p_i^2/2m + n\hbar\omega$.

We can see in equation (16) the cross-section of the scattering by radiation and the cross-section of the scattering from the potential V ; in addition, there appear mixed radiation-potential scattering terms; for $n = 0$ we get from equation (16) the elastic Born scattering in the field of the potential V affected by radiation, due to the presence of the function J_0 . The argument of the Bessel function in equation (16) varies rapidly over the integration domain; therefore, we may use the asymptotic expression for the Bessel function; doing so, we can see easy that the \mathbf{r} -integration is non-vanishing for $\mathbf{p} \simeq e\mathbf{E}_0/\omega$, *i.e.* the momentum is trasferred along the direction of the electric field, as expected; this is the momentum conservation. Making use of this relation and the energy conservation we get $p_{fn} \simeq p_i(1 + n\hbar\omega/2E_i)$ (for $n\hbar\omega \ll E_i$) and

$$-1 < \cos \theta_n = \frac{1 - e\mathbf{p}_i\mathbf{E}_0/p_i^2\omega}{1 + n\hbar\omega/2E_i} < 1 , \quad (17)$$

where θ_n is the scattering angle. For $n > 0$ (photon absorption) the scattering angle θ_n increases, while for $n < 0$ (photon emission) the scattering angle decreases, with respect to the elastic scattering angle θ_0 . Equation (17) indicates that there exists a limitation which can be written as $-n\hbar\omega/2E_i < (eA_0 \cos \alpha/2E_i)(v/c) < 2 + n\hbar\omega/2E_i$, where α is the angle the initial momentum makes with the electric field; for very low incident energies the scattering occurs only for angles close to the right angle between the incident momentum and the electric field. Finally, we note that the order of magnitude of the cross-section of the scattering due to radiation is $\simeq d^2[n\hbar\omega/(\hbar^2/md^2)]^2(\hbar c/eA_0d)$, where $(\hbar c/eA_0d) \ll 1$ and d is the dimension of the region where the radiation is confined to. We can see from equation (16) that the cross-section due to radiation may acquire large values, as a consequence of the large dimension of the region containing radiation; the cross-section increases with increasing n , *i.e.* for large scattering angles, where, however, the scattering maxima coalesce.

The above calculations are done for a sufficiently long laser pulse. In practice, the pulse has a finite duration τ and a repetition time Δt . In these conditions the expansion given by equation (13) remains valid, but the function $\delta(\Delta E)$ in the scattering amplitude a_{fi} (equation (14)), where $\Delta E = E_f - E_i - n\hbar\omega$, is replaced by the function $\zeta(\Delta E) = e^{\frac{i}{\hbar}\Delta E t_i} \sin \alpha \Delta E / \pi \Delta E$, where $\alpha = \tau/2\hbar$ and t_i denotes the time of the pulse (the pulse lasts from $t_i - \tau/2$ to $t_i + \tau/2$). For large α the function $\zeta(\Delta E)$ has a maximum for $\Delta E = 0$ and extends approximately over a bandwidth $\delta E \simeq \pi/\alpha = 2\pi\hbar/\tau$. Since, usually τ is much longer than the radiation period $T = 2\pi/\omega$, the energy separation δE is much smaller than the radiation quanta of energy $\hbar\omega$. It folows that the functions $\zeta(\Delta E)$ for different n can be viewed as being well separated. In these conditions the cross-section $d\sigma_n$ (equation (16)) preserves its form, except that it is multiplied by the reduction factor $\tau/(\tau + \Delta t)$.

In conclusion, it is shown in this paper that non-relativistic charged particles may suffer scattering as a result of their interaction with the electromagnetic radiation in the focal region of laser beams. The cross-section of this scattering process is computed in this paper for a single-mode radiation with linear polarization. As expected, the cross-section exhibits maxima at certain scattering (diffraction) angles θ_n , as given by equation (17), determined by the energy and momentum conservation in multiple-photon exchange processes. The calculations can be immediately extended to any polarization; for realistic laser beams, or for multi-mode radiation, we should take into account the particular beam shape and the amplitude and frequency fluctuations. For convenience, we included also the scattering from a potential placed in the radiation field, in the Born approximation. The cross-section of the potential scattering is modified by the presence of the radiation, because the scattering states are dressed by radiation. The cross-section is reduced for high-intensity radiation (preserving the non-relativistic character of the particle motion) and, similarly, the multi-photon scattering is reduced for high energy of the particle, when the process reduces to elastic scattering in the forward direction. In contrast with previous studies, where the scattering process is immersed in the radiation field, in the scattering process addressed in this paper the radiation field is confined to a finite region.

Acknowledgements. The author is indebted to the members of the Laboratory of Theoretical Physics at Magurele-Bucharest for many fruitful discussions. This work has been supported by the Scientific Research Agency of the Romanian Government through Grants 04-ELI / 2016 (Program 5/5.1/ELI-RO), PN 16 42 01 01 / 2016 and PN (ELI) 16 42 01 05 / 2016

References

- [1] N. M. Kroll and K. M. Watson, "Charged-particle scattering in the presence of a strong electromagnetic wave", *Phys. Rev.* **A8** 804-809 (1973).
- [2] N. K. Rahman, "Effect of intense electromagnetic wave in charged-particle scattering", *Phys. Rev.* **A10** 440-441 (1974).
- [3] J. Bergou, "Wavefunctions of a free electron in an external field and their application in intense field interactions. I. Non-relativistic treatment", *J. Phys.* **A13** 2817-2822 (1980).
- [4] J. Bergou and S. Varro, "Wavefunctions of a free electron in an external field and their application in intense field interactions. II. Relativistic treatment", *J. Phys.* **A13** 2823-2837 (1980).
- [5] A. F. Klinskikh and L. P. Rapoport, "Quasi-classical theory of potential scattering of an electron in the field of a laser wave", *Sov. Phys.-JETP* **66** 53-57 (1987) (*ZhETF* **93** 95-103 (1987)).
- [6] N. J. Mason, "Laser-assisted electron-atom collisions", *Reps. Progr. Phys.* **56** 1275-1346 (1993).
- [7] B. Wallbark and J. K. Holmes, "Laser-assisted elastic electron-atom collisions: low electron energy and small scattering angle", *J. Phys. B: At. Mol. Opt. Phys.* **27** 1221-1231 (1994)
- [8] B. Wallbark and J. K. Holmes, "Differential cross sections for laser-assisted elastic electron scattering from argon", *J. Phys. B: At. Mol. Opt. Phys.* **27** 5405-5418 (1994)

- [9] L. B. Madsen and K. Taulbjerg, "Theory of laser-assisted electron-atom scattering: beyond the Kroll-Watson approximation, J. Phys. B: At. Mol. Opt. Phys. **28** 5327-5342 (1995)
- [10] I. F. Barna and S. Varro, "Laser-assisted proton collision on light nuclei at moderate energies", Laser&Part. Beams **33** 299-305 (2015).
- [11] Y. I. Salamin, S. X. Hu, K. Z. Hatsagortsyan and C. H. Keitel, "Relativistic high-power laser-matter interaction", Phys. Reps. **427** 41-155 (2006).
- [12] M. Marklund and P. K. Shukla, "Nonlinear collective effects in photon-photon and photon-plasma interactions", Revs. Mod. Phys. **78** 591-640 (2006).
- [13] H. K. Avetissian, *Relativistic Nonlinear Electrodynamics. Interaction of Charged Particles with Strong and Super Strong Laser Fields*, Springer Series in Optical Sciences **120**, ed. W. T. Rhodes, Springer, New York (2006).
- [14] V. Yanovski, V. Chvykov, G. Kalinchenko, P. Rousseau, T. Planchon, T. Matsuoka, A. Maksimchuk, J. Nees, G. Cheriaux, G. Mourou and K. Krushelnik, "Ultra-high intensity-300-Tw laser at 0.1Hz repetition rate", Opt. Express **16** 2109-2114 (2008).
- [15] F. Ehlotzky, K. Krajewska and J.Z. kaminski, "Fundamental processes of quantum electrodynamics in laser fields of relativistic power", Reps. Progr. Phys. **72b** 046401 (1-32) (2009).
- [16] A. V. Korzhimanov, A. A. Gonoskov, E. A. Khazanov and A. M. Sergeev, "Horizons of petawatt laser technology", Phys. Usp. **54** 9-28 (2011).
- [17] A. di Piazza, C. Muller, K. Z. Hatsagortsyan and C. H. Keitel, "Extremely high-intensity laser interactions with fundamental quantum systems", Revs. Mod. Phys. **84** 1177- 1228 (2012).
- [18] V. Malka, "Laser plasma accelerators", Phys. Plasmas **19** 055501 (2012).
- [19] P. L. Kapitza and P. A. M. Dirac, "The reflection of electrons from standing light waves", Proc. Cambridge Phil. Soc. **29** 297-300 (1933).
- [20] M. Goeppert-Mayer, "Uber Elementarakte mit zwei Quantensprungen", Ann. Physik **401** 273-294 (1931).
- [21] W. C. Henneberger, "Perturbation method for atoms in intense light beams", Phys. Rev. Lett. **21** 838-840 (1968).
- [22] H. A. Kramers, "Non-relativistic quantum-electrodynamics and the correspondence principle", Solvay Congress 1948, in *Collected Scientific Papers*, North-Holland, Amsterdam (1956), p. 865.
- [23] N. G. van Kampen, "Contribution to the quantum theory of light scattering", Kgl. Danske Videnskab. Selskab, Mat.-fys. Medd. **26** (15) (1951).