Journal of Theoretical Physics

Founded and Edited by M. Apostol

ISSN 1453-4428

Classical interaction of the electromagnetic radiation with two -level polarizable matter

M. Apostol

Department of Theoretical Physics, Institute of Atomic Physics, Magurele-Bucharest MG-6, POBox MG-35, Romania email: apoma@theory.nipne.ro

Abstract

The interaction of the classical electromagnetic field with an ensemble of polarizable, identical, atomic particles with two energy levels is investigated, and the coupled non-linear equations of motion for the polarization field and the amplitudes of the level occupancies are solved by a perturbation-theoretical method. A small coupling constant is identified, and the solution is represented as a power series in this coupling constant. Explicit results are given for the leading contributions to the solution. In particular, it is shown that an external electromagnetic field may induce a lasing effect in such an ensemble of particles, by populating the (initially empty) upper level.

Key words: matter interacting with radiation, polarization, lasers, perturbation theory, non-linear equations

PACS: 41.20.-q; 42.55.Ah; 42.50.Ct; 42.50.Pq

The interaction of the classical electromagnetic radiation with an ensemble of polarizable, identical, atomic particles with two energy levels is the core of the "semi-classical theory" of the laser (see, for instance, Refs. [1]-[3]). The problem has been extensively investigated, by various approaches and from many angles.[4]-[17] Usually, the equations of motion for the electromagnetic field and the occupancies of the two levels are solved by means of some approximations which, among other particular assumptions, discard the fast oscillating terms. However, such terms may bring relevant contributions in the stationary regime. It is generally believed that an exact solution of the coupled, non-linear equations of the semi-classical theory of the laser would be impossible (see, for instance, Ref. [2], p. 459, Ref. [3], p. 98). We present here a fully computable solution, represented as a power series in a (small) coupling constant λ , and give explicit results for the polarization field, occupancy numbers and energy in the lowest, most relevant orders of λ , in the ensemble of particles, driven by the external field which can populate the (initially empty) upper level.

We consider a uniform distribution of polarizable, identical particles, each with two quantum energy levels $\varepsilon_{0,1}$, subjected to an external electromagnetic field and to their own polarization field. The ensemble of particles exhibits a fluctuating current density $\mathbf{j}(\mathbf{r},t)$, and a polarization $\mathbf{P}(\mathbf{r},t)$, related by $\mathbf{j}(\mathbf{r},t) = \partial \mathbf{P}(\mathbf{r},t)/\partial t$, which, in turn, give rise to a polarization field, according to the well-known wave equations with sources

$$\frac{1}{c^2}\frac{\partial^2 \mathbf{A}}{\partial t^2} - \Delta \mathbf{A} = \frac{4\pi}{c}\mathbf{j} \ , \ \frac{1}{c^2}\frac{\partial^2 \mathbf{E}}{\partial t^2} - \Delta \mathbf{E} = -\frac{4\pi}{c^2}\frac{\partial^2 \mathbf{P}}{\partial t^2} \ , \tag{1}$$

where **A** is the vector potential and $\mathbf{E} = -(1/c)\partial \mathbf{A}/\partial t$ is the polarization electric field (we assume a transverse radiation field). We take only one polarization, oriented along one coordinate axis, and look for a separable solution of the form $E(\mathbf{r},t) = E(t)\chi(\mathbf{r})$, $P(\mathbf{r},t) = P(t)\chi(\mathbf{r})$, where $\chi(\mathbf{r})$ is an eigenfunction of the laplacian, $\Delta\chi(\mathbf{r}) = -\kappa^2\chi(\mathbf{r})$, κ being a constant. With the notation $\omega_0^2 = c^2\kappa^2$, the second equation (1) becomes

$$\ddot{E}(t) + \omega_0^2 E(t) = -4\pi \frac{\partial^2 P(t)}{\partial t^2} .$$
⁽²⁾

We envisage a classical polarization field E; consequently, the source in the *rhs* of equation (2) can be written as $4\pi n \langle \partial^2 p / \partial t^2 \rangle$, where p is the dipole momentum of a particle and the brackets denote the quantum average; the spatial average is taken into account by the (uniform) density n of the ensemble of particles. We take $\langle \partial^2 p / \partial t^2 \rangle = -\omega_1^2 \langle p \rangle$, where $\hbar \omega_1 = \varepsilon_1 - \varepsilon_0$. Equation (2) can then be written as

$$\ddot{E}(t) + \omega_0^2 E(t) = 4\pi n \omega_1^2 \langle p \rangle \quad . \tag{3}$$

In general, $\langle \partial^2 p / \partial t^2 \rangle$ depends on the internal dynamics of the particles, and can be kept as such in equation (3), or it may be expressed in terms of other conventional parameters. We note also that the electric field source, given generally by $\partial j(t) / \partial t$, may not originate only in orbital currents (as we assumed here), but it may have also other origins, like the spin, for instance.

The two quantum states $\varphi_{0,1}$ are defined by the free hamiltonian H_0 of the internal degrees of freedom of each individual particle, $H_0\varphi_{0,1} = \varepsilon_{0,1}\varphi_{0,1}$. The interaction hamiltonian for one particle placed at **r** is given by

$$H_{int} = -p\chi(\mathbf{r}) \left[E_0(\mathbf{r}, t) + E(\mathbf{r}, t) \right] = -pE_t(t)\chi^2(\mathbf{r}) \quad , \tag{4}$$

where the external field E_0 has been introduced, as well as the total field $E_t = E_0 + E$. We assume the fields and the (orthogonalized) eigenfunctions real. The spatial average of equation (4) gives an interaction hamiltonian

$$(H_{int})_{av} = -pE_t(t) . (5)$$

The interacting state $\varphi = c_0 \varphi_0 + c_1 \varphi_1$ is a superposition of the two free states $\varphi_{0,1}$, with coefficients $c_{0,1}$ satisfying the Schrödinger equation

$$i\hbar\frac{\partial c_0}{\partial t} = \varepsilon_0 c_0 - p_{01} E_t c_1 ,$$

$$i\hbar\frac{\partial c_1}{\partial t} = \varepsilon_1 c_1 - p_{01}^* E_t c_0 .$$
(6)

The quantum average of the current is given by

$$\langle p \rangle = p_{01}c_0^*c_1 + p_{01}^*c_1^*c_0 \quad , \tag{7}$$

where we have assumed $p_{00} = p_{11} = 0$, as for stationary states. Moreover, we assume for simplicity $p_{01} = p_{01}^* = p$. We set $\varepsilon_0 = 0$ and introduce the parameter

$$x(t) = \frac{2p}{\hbar\omega_1} E_t(t) \quad , \tag{8}$$

so that equations (6) become

$$i\frac{\partial c_0}{\partial t} = -\frac{1}{2}\omega_1 x(t)c_1 , \ i\frac{\partial c_1}{\partial t} = \omega_1 c_1 - \frac{1}{2}\omega_1 x(t)c_1$$
(9)

and equation (3) can be written as

$$\ddot{E}(t) + \omega_0^2 E(t) = 4\pi n \omega_1^2 p \left(c_0^* c_1 + c_1^* c_0 \right) .$$
(10)

In equation (8) we may recognize the well-known Rabi "frequency" pE_t/\hbar . Usually, the system of equations (9) is transformed into a system of equations for the occupancies $|c_{0,1}|^2$ and the associated matrix density.[2, 3] We adopt a different route, and focus on the system of equations (9) for the occupancy amplitudes $c_{0,1}$.

The system of equations (9) can be solved formally with $c_{0,1} = C_{0,1}e^{i\theta}$; we get immediately $\dot{C}_{0,1} = 0$ and

$$c_{0} = C_{0}e^{i\theta_{0}} - fC_{1}e^{i\theta_{1}} , \ c_{1} = fC_{0}e^{i\theta_{0}} + C_{1}e^{i\theta_{1}} ,$$

$$\dot{\theta}_{0,1} = \frac{1}{2}\omega_{1} \left(-1 \pm \sqrt{x^{2}(t) + 1}\right) ,$$
(11)

where

$$f(t) = \frac{x(t)}{\sqrt{x^2(t) + 1} + 1}$$
 (12)

The coefficients $C_{0,1}$ are determined by requiring the initial values of the occupancy numbers $|c_{0,1}(t=0)|^2$ be equal with $n_{0,1}$ $(n_0 + n_1 = 1)$. We get the amplitudes

$$C_{0,1} = \frac{1}{1 + f^2(t)} \left[\sqrt{n_{0,1}} \pm f(t) \sqrt{n_{1,0}} \right]$$
(13)

and the occupancy numbers

$$|c_{0,1}|^2 = n_{0,1} \pm \frac{1}{2} \frac{x(t)}{x^2(t) + 1} \left[2\sqrt{n_0 n_1} - x(t)(n_0 - n_1) \right] \left[1 - \cos(\theta_0 - \theta_1) \right] , \qquad (14)$$

where the phase difference $\theta_0 - \theta_1$ is given by

$$\Delta \theta = \theta_0 - \theta_1 = \omega_1 \int_0^t dt \sqrt{x^2(t) + 1} .$$
 (15)

The oscillations of the occupancies given by equation (14) are reminiscent of the well-known Rabi oscillations, exhibited, for instance, by the Jaynes-Cummings model (see, for instance, Refs. [4, 15]). We take the time averages of all the relevant quantities given above. We can see, by equations (11), that the energy levels $\varepsilon_{0,1}$ are changed by interaction into the mean values of $\hbar \dot{\theta}_{0,1}$, and, in addition, the interaction mixes up the two states, as expected. We can see also that the mean values of the coefficients $C_{0,1}$, as well as the mean values of the coefficients $fC_{0,1}$ entering equations (11), are constants, as it is required by a stationary solution; it becomes apparent that $n_{0,1}$ are constants of integration.

From equations (11)-(13) we get

$$c_0^* c_1 + c_1^* c_0 = \frac{1}{x^2 + 1} \left\{ x \left[2x \sqrt{n_0 n_1} + n_0 - n_1 \right] + \left[2\sqrt{n_0 n_1} - x(n_0 - n_1) \right] \cos \Delta \theta \right\} ,$$
(16)

which can be inserted into equation (10); we can add the external field E_0 , which satisfies the free wave equation $\ddot{E}_0 + \omega_0^2 E_0 = 0$, such that equation (10) becomes

$$\ddot{x} + \omega_0^2 x = \lambda^2 \omega_1^2 \frac{1}{x^2 + 1} \left\{ x \left[2x \sqrt{n_0 n_1} + n_0 - n_1 \right] + \left[2\sqrt{n_0 n_1} - x(n_0 - n_1) \right] \cos \Delta\theta \right\} ,$$
(17)

where $\lambda^2 = 8\pi np^2/\hbar\omega_1$. We note that $x = \lambda E_t/e$, where $e = \sqrt{2\pi n\hbar\omega_1}$ is a characteristic electric field.

Equation (17) is a non-linear (integro-differential) equation. We assume $\lambda \ll 1$, and seek the solution as a power series in λ ,

$$x = \lambda x_0 + \lambda^2 x_1 + \lambda^3 x_2 + \dots, \qquad (18)$$

where $x_0 = B \cos \widetilde{\omega}_0 t$, $B = E_0/e$ and $\widetilde{\omega}_0$ remains to be determined. We get straightforwardly

$$x_{1} = 2\sqrt{n_{0}n_{1}} \frac{\omega_{1}^{2}}{\omega_{0}^{2}-\omega_{1}^{2}} \cos\widetilde{\omega}_{1}t ,$$

$$\frac{1}{2}(n_{0}-n_{1})B\left[\frac{\omega_{1}}{2\omega_{0}+\omega_{1}}\cos(\widetilde{\omega}_{0}+\widetilde{\omega}_{1})t - \frac{\omega_{1}}{2\omega_{0}-\omega_{1}}\cos(\widetilde{\omega}_{0}-\widetilde{\omega}_{1})t\right] ,$$
(19)

where

 $x_2 =$

$$\widetilde{\omega}_{0} = \omega_{0} - \lambda^{2} \frac{\omega_{1}^{2}}{2\omega_{0}} (n_{0} - n_{1}) , \ \widetilde{\omega}_{1} = \omega_{1} \left(1 + \frac{1}{4} \lambda^{2} B^{2} \right) , \qquad (20)$$

for $\omega_1 \neq \omega_0, \pm 2\omega_0$. These restrictions can be related to the parametric resonances $2\omega_0 \simeq n\omega_1$ (where $n \neq 0$ is any integer), occurring for an associated Mathieu equation which is a close representation of the linearized form of equation (17) for $n_1 = 0$ (though not a fully correct approximation to equation (17)).[18] Leaving aside the (weak) frequency renormalization, the resonances exhibited by equations (19) are in fact what we may expect from a non-linear oscillator with the basic frequency ω_0 subjected to an external force of frequency ω_1 . As it is well known, such an oscillator exhibits the combined-frequency phenomenon, as reflected in the occurrence of frequencies of the form $\omega_0 \pm \omega_1$ and denominators $2\omega_0 \pm \omega_1$, etc (arising from terms like $\omega_0^2 - (\omega_0 \pm \omega_1)^2$).

We can see that the interaction renormalizes both the field frequency ω_0 and the characteristic frequency ω_1 of the ensemble of particles. The term x_1 represents the oscillations of the ensemble of particles (for $n_{0,1} \neq 0$); the effect of the external field appears only in the next order (the term x_2), with combined frequencies $\tilde{\omega}_0 \pm \tilde{\omega}_1$. For $n_{1,0} = 0$ the polarization process is governed entirely by the external field, as expected (and the constraint $\omega_0 \neq \omega_1$ is removed).

Having known the parameter x(t), the mean values (time averages) of all the relevant quantities can be computed, as given by equations (11)-(15). We get, for instance, the frequencies

$$\Omega_0 = \overline{\dot{\theta}}_0 = \frac{1}{8} \lambda^2 \omega_1 B^2 , \ \Omega_1 = \overline{\dot{\theta}}_1 = -\omega_1 - \frac{1}{8} \lambda^2 \omega_1 B^2$$
(21)

and the mean occupancies

$$\overline{|c_{0,1}|^2} = n_{0,1} \mp \lambda^2 n_0 n_1 \frac{\omega_1^2}{\omega_0^2 - \omega_1^2} \mp \frac{1}{4} \lambda^2 (n_0 - n_1) B^2 .$$
(22)

One can see that the external field can pump, or deplete, the upper level, depending on the parameters $n_{0,1}$ and $\omega_{0,1}$. Particularly interesting is the case $n_1 = 0$ (corresponding to an upper level which is empty at the initial moment t = 0). In this case, the occupancy of the upper level is given by

$$\overline{|c_1|^2} = \frac{1}{4}\lambda^2 B^2 = \frac{1}{4e^2}\lambda^2 E_0^2 = \left(\frac{pE_0}{\hbar\omega_1}\right)^2 ; \qquad (23)$$

the external field leads to a macroscopic occupation of this level. The release of the corresponding energy $E_s = \hbar \omega_1 |c_1|^2$ (per particle) is a lasing effect, driven by the external field. We note in equation (23) the occurrence of the Rabi frequency pE_0/\hbar .

The polarization can be computed by making use of the solution x(t) given here (equation (19)) in equations (7) and (16). Within this approximation, the polarization is a complicate function, involving quadratic dependence on the strength of the external field, frequency doubling, combined frequencies, etc, as expected for a non-linear equation. We collect here a few relevant terms:

$$c_0^* c_1 + c_1^* c_0 = 2\sqrt{n_0 n_1} \cos \widetilde{\omega}_1 t + \lambda B(n_0 - n_1)(1 - \cos \widetilde{\omega}_1 t) \cos \widetilde{\omega}_0 t +$$

$$+ 2\lambda^2 \sqrt{n_0 n_1} \left[B^2 \cos^2 \widetilde{\omega}_0 t - (n_0 - n_1) \frac{\omega_1^2}{\omega_0^2 - \omega_1^2} \cos^2 \widetilde{\omega}_1 t \right] -$$

$$+ \lambda^3 \frac{\omega_1 B^2}{16\omega_0} (n_0 - n_1) \sin \widetilde{\omega}_0 t \sin \widetilde{\omega}_1 t .$$
(24)

The mean value of the polarization is given by

$$\overline{P} = n \overline{p \left(c_0 c_1^* + c_1 c_0^* \right)} = \lambda^2 n p \sqrt{n_0 n_1} \left[B^2 - (n_0 - n_1) \frac{\omega_1^2}{\omega_0^2 - \omega_1^2} \right]$$
(25)

where the quadratic dependence on the external field is to be noted. It is worth noting that $\overline{P} = 0$ for $n_{0,1} = 0$. The $\tilde{\omega}_0$ -component of the polarization gives the permittivity $\kappa = (2np^2/\hbar\omega_1)(n_0 - n_1)$ and the particle polarizability $\alpha = 2p^2/\hbar\omega_1$ (for $n_1 = 0$). We can see that the coupling constant $\lambda = \sqrt{8\pi np^2/\hbar\omega_1}$ can be related to the polarizability α through $\lambda^2 = 4\pi n\alpha$. It follows that we are justified in assuming $\lambda \ll 1$ as far as the polarizability per unit volume is small, which is a usual situation.

By equation (18), the total electric field can be represented as a $E_t = e(x_0 + \lambda x_1 + \lambda^2 x_2 + ...)$. Hence, we can obtain the vector potential $A_t(t) = -c \int_0^t dt E_t$ and the magnetic field $\mathbf{H} = curl \mathbf{A}_t(\mathbf{r}, t)$ (where we take into account the transversality condition $div \mathbf{A}_t = 0$). It is then easy to compute the mean value of the total field energy density (per particle) E_t^t . We get

$$E_f^t = \frac{1}{8\pi n} E_0^2 + \frac{\lambda^2}{4\pi n} \left[n_0 n_1 \frac{\omega_1^2(\omega_0^2 + \omega_1^2)}{(\omega_0^2 - \omega_1^2)^2} e^2 + (n_0 - n_1) \frac{\omega_1^2}{4\omega_0^2} E_0^2 \right] .$$
(26)

From equations (5), (7) and (16) we can compute the interaction energy (per particle)

$$E_{int} = -\langle p \rangle E_t = -\frac{1}{2}\hbar\omega_1 x \left(c_0^* c_1 + c_1^* c_0 \right)$$
(27)

and its mean value

$$\overline{E}_{int} = -\lambda^2 \hbar \omega_1 \left[n_0 n_1 \frac{\omega_1^2}{\omega_0^2 - \omega_1^2} + \frac{1}{4} (n_0 - n_1) B^2 \right] .$$
(28)

As it was said above, the energy of the ensemble of particles (per particle) is given by

$$E_{s} = \hbar\omega_{1} \overline{|c_{1}|^{2}} = \hbar\omega_{1} \left[n_{1} + \lambda^{2} n_{0} n_{1} \frac{\omega_{1}^{2}}{\omega_{0}^{2} - \omega_{1}^{2}} + \frac{1}{4} \lambda^{2} (n_{0} - n_{1}) B^{2} \right] .$$
(29)

Particularly interesting are these equations for $n_1 = 0$:

$$E_f^t = E_f^0 \left(1 + \lambda^2 \frac{\omega_1^2}{2\omega_0^2} \right) , \qquad (30)$$
$$E_s = -\overline{E}_{int} = \frac{1}{4} \hbar \omega_1 \lambda^2 B^2 = \lambda^2 E_f^0 ,$$

where $E_f^0 = E_0^2/8\pi n$ is the energy (per particle) of the (bare) external field. One can see that the total energy $E_t = E_f^t + E_s + \overline{E}_{int}$ reduces to the total field energy E_f^t , the polarization energy (E_s) being entirely compensated by the interaction energy (\overline{E}_{int}) , as expected. The efficiency quotient of this lasing process is $\simeq \lambda^2$.

In conclusion, after setting up the equations of motion for the polarization field and the amplitudes of the level occupancy for an ensemble of identical, polarizable particles with two energy levels interacting with a classical electromagnetic radiation, we have identified a small coupling constant λ which allows the solution of these coupled non-linear equations to be obtained by a theoreticalperturbation method. The solution is represented as a power series in λ . The coupling constant λ is related to the polarizaton $n\alpha$ per unit volume of the ensemble of particles, $\lambda = \sqrt{4\pi n\alpha}$. Explicit results have been given for the leading contributions to the polarization field, occupancy numbers and energy in the presence of an external field. It was shown that an external field can induce a lasing effect, by pumping the upper energy level which was empty at the initial moment of time.

Finally, it is worthwhile commenting upon the absence of the external field in the problem presented in this paper. The equations of motion (3) and (6) have a completely different solution in this case. These equations can be solved exactly, and the solution, which is independent of time and exists only for λ exceeding a critical value, is non-analytic in the coupling constant λ . For a strong enough coupling the ensemble of particles is unstable, and it undergoes a phase transition to a state known as a super-radiance state. Beside such a criticality condition upon the coupling constant, for finite temperatures the super-radiance (second-order) phase transition occurs below a certain critical temperature. The exact solution was obtained in a previous paper, [19] where the super-radiance transition was also characterized, under the assumption of a coherent interaction of matter with radiation. It was shown there that the two energy levels and the photon states are macroscopically occupied, leading to a spatially long-range ordered state which involves the quantum phases of the internal motion of the particles. The transition process is initiated, and carried on, for $\omega_0 = \omega_1$, as corresponding to quantum transitions. In the classical case the condition of coherent interaction is implicitly included. The solution of the classical equations of motion (3) and (6) in the absence of the external field is identical with the solution obtained in Ref. [19] within the coherent interaction theory.

References

- [1] M. Sargent, M. O. Scully and W. E. Lamb, *Laser Physics*, Addison-Wesley, Reading (1974).
- [2] W. H. Louisell, Quantum Statistical Properties of Radiation, Wiley, NY (1973).
- [3] H. Haken, Laser Theory, in Encyclopedia of Physics, vol. XXV/2c, S. Flugge ed., Springer, Berlin (1970).
- [4] E. T. Jaynes and F. W. Cummings, Comparison of quantum and semiclassical radiation theories with application to the beam maser, Proc. IEEE **51** 89-109 (1963).
- S. Stenholm, Quantum theory of electromagnetic fields interacting with atoms and molecules, Phys. Reps. C6 1-121 (1973).
- [6] B. R. Mollow, Pure-state analysis of resonant light scattering: radiative damping, saturation and multiphoton effects, Phys. Rev. A12 1919-1943 (1975).

- [7] C. Cohen-Tannoudji, in Frontiers in Laser Spectroscopy, vol. 1, Les Houches 1975, eds. R. Balian, S. Haroche and S. Liberman, North-Holland, NY (1977).
- [8] L. Allen, J. H. Eberly, *Optical Resonance in Two-Level Atoms*, Dover, NY (1987).
- [9] R. Loudon, The Quantum Theory of Light, Oxford University, NY (1983).
- [10] J. Dalibard and C. Cohen-Tannoudji, Dressed-atom approach to atomic motion in laser light: the dipole force revisited, J. Opt. Soc. Am. B2 1707-1720 (1985).
- [11] B. W. Shore, The Theory of Coherent Atomic Excitations, Wiley, NY (1990).
- [12] C. Cohen-Tannoudji, J. Dupont-Roc and G. Grynberg, Atom-Photon Interactions: Basic Processes and Applications, Wiley, NY (1992).
- [13] O. Kocharovskaya, Amplification and lasing without inversion, Phys. Reps. 219 175-190 (1992).
- [14] H. C. Fogedby, Field-theoretical derivation of the master equation in quantum optics, Phys. Rev. A47 4364-4383 (1993).
- [15] B. W. Shore and P. L. Knight, The Jaynes-Cummings model, J. Mod. Optics 40 1195-1238 (1993).
- [16] K. Bergmann, H. Theuer and B. W. Shore, Coherent population transfer among quantum states of atoms and molecules, Revs. Mod. Phys. 70 1003-1025 (1998).
- [17] M. Fleischhauer, A. Imamoglu and J. P. Marangos, Electromagnetically induced transparency: Optics in coherent media, Revs. Mod. Phys. 77 633-673 (2005).
- [18] See, for instance, E. T. Whittaker and G. N. Watson, A Course of Modern Analysis, Cambridge University Press, Cambridge (2004).
- [19] M. Apostol, Coherence domains in mater interacting with radiation, Phys. Lett. A373 379-384 (2009) and references therein.

ⓒ J. Theor. Phys. 2010, apoma@theor1.theory.nipne.ro