

# Randomness effects on modulational instability of a discrete self-trapping equation

Anca Vişinescu, D. Grecu

*Department of Theoretical Physics*

*National Institute for Physics and Nuclear Engineering*

*"Horia Hulubei"*

*P.O.Box MG-6, Măgurele, Bucharest, Romania*

*e-mail: avisin@theor1.theory.nipne.ro*

*dgrecu@theor1.theory.nipne.ro*

## Abstract

The discrete self-trapping equation (DST) represents an useful model for several properties of one-dimensional nonlinear molecular crystals. The modulational instability of DST equation is discussed from a statistical point of view, considering the oscillator amplitude as a random variable. A kinetic equation for the two-point correlation function is written down, and its linear stability is studied. Both a Gaussian and a Lorentzian form for the initial unperturbed wave spectrum are discussed. Comparison with the continuum limit (NLS equation) is done.

## 1 Introduction

The discrete self-trapping (DST) equation

$$i \frac{da_n}{dt} - \omega_0 a_n + \lambda(a_{n+1} + a_{n-1}) + \mu |a_n|^2 a_n = 0 \quad (1)$$

is a typical equation for a system of harmonically coupled nonlinear oscillations [1], [2] relevant for several physical problems. We mention here only Davydov's model of energy transport in  $\alpha$ -helix structures in proteins [3], [4], [2], where (1) appears as a certain approximation of the model. In (1)  $a_n$  is the complex classical dimensionless amplitude of the oscillator of frequency  $\omega_0$  in the  $n$ -th molecule, and  $\lambda, \mu$  (of dimension of frequency) are the coupling constants between nearest neighbour oscillators and the one-site nonlinearity respectively. It is well known that depending upon of the parameters and the chosen initial condition the equation (1) can lead either to self-trapping (i.e. local modes or solitons), or to chaos, or to a mixture of the above two behaviours [1], [2], [5]. Instead of (1) we shall consider the equation

$$i \frac{da_n}{dt} + \lambda(a_{n+1} + a_{n-1}) + \mu|a_n|^2 a_n = 0 \quad (2)$$

which is obtained if  $a_n \rightarrow a_n e^{-i\omega_0 t}$ . This equation admits plane wave solutions with constant amplitude

$$a_n = a e^{i(kn - \omega t)}$$

(the lattice constant is taken equal with unity) but with an amplitude depending dispersion relation

$$\omega(k) = -2\lambda \cos k - \mu|a|^2$$

This is a Stokes wave solution and it is well known to be unstable at small modulation of the amplitude (Benjamin-Feir or modulational instability) [6]-[8]. The aim of this note is to study the modulational instability of equation (2) from a statistical point of view considering  $a_n$  as a random variable. In doing this we shall follow the procedure used by several authors to discuss the effects of randomness on the stability of weakly nonlinear waves, especially in hydrodynamics [9], [10].

In the next section a kinetic equation for a two-point correlation function will be obtained. Using a Wigner-Moyal transform the equation is written in a mixed configuration-wave vector space. The linear stability around a homogeneous basic solution is discussed in section 3. An integral stability equation is derived, very similar with the dispersion relation of the linearized Vlasov equation in ionized plasmas. Two forms for the spectrum of the initial unperturbed condition will be considered, namely a Gaussian and a Lorentzian form and in the limit of vanishingly small bands widths the

increment of the modulational instability is calculated. Comparison with the continuum limit, when (1) transforms into the nonlinear Schrödinger equation is done. Few concluding remarks are also presented.

## 2 Kinetic equation for two-point correlation function

Introducing the displacement operator by  $a_{n\pm 1} = e^{\frac{\partial}{\partial n}} a_n$  the equation (2) becomes

$$i \frac{\partial a_n}{\partial t} + 2\lambda \cosh \frac{\partial}{\partial n} a_n + \mu |a_n|^2 a_n = 0. \quad (3)$$

In order to find a kinetic equation we write (3) for  $n = n_1$ , multiply it by  $a_{n_2}^*$ , add it to the complex conjugated of (3) for  $n = n_2$  multiplied by  $a_{n_1}$  and finally take an ensemble average. One obtains

$$i \frac{\partial}{\partial t} \langle a_{n_1} a_{n_2}^* \rangle + 2\lambda (\cosh \frac{\partial}{\partial n_1} - \cosh \frac{\partial}{\partial n_2}) \langle a_{n_1} a_{n_2}^* \rangle + \mu (\langle a_{n_1} a_{n_1}^* a_{n_1} a_{n_2}^* \rangle - \langle a_{n_2} a_{n_2}^* a_{n_1} a_{n_1}^* \rangle) = 0 \quad (4)$$

which beside the two point correlation function  $\rho(n_1, n_2, t) = \langle a_{n_1}(t) a_{n_2}^*(t) \rangle$  contains also four-point correlation functions. If  $a_n$  corresponds to a Gaussian process, and this property is retained during the evolution, a four-point correlation function factorizes exactly in products of two-point correlation functions [11]

$$\langle a_{n_1} a_{n_1}^* a_{n_1} a_{n_2}^* \rangle = 2 \langle a_{n_1} a_{n_2}^* \rangle \langle a_{n_1} a_{n_1}^* \rangle = 2\rho(n_1, n_2) \bar{a}^2(n_1) \quad (5)$$

where  $\bar{a}^2(n) = \langle a_n a_n^* \rangle$  is the ensemble average of the mean square amplitude. Although the factorization (5) is true only for a Gaussian process we shall assume to be at least approximately valid also for processes slightly different from a Gaussian one, and it represents the main approximation of the present analysis.

It is convenient to use a Wigner-Moyal transform [12]. One introduce the new variables

$$M = \frac{n_1 + n_2}{2}, \quad m = n_1 - n_2. \quad (6)$$

Then

$$\cosh \frac{\partial}{\partial n_1} - \cosh \frac{\partial}{\partial n_2} = 2 \sinh \frac{1}{2} \frac{\partial}{\partial M} \sinh \frac{\partial}{\partial m}$$

and equation (4) becomes

$$i \frac{\partial \rho}{\partial t} + 4\lambda \sinh \frac{1}{2} \frac{\partial}{\partial M} \sinh \frac{\partial \rho}{\partial m} + 2\mu (\bar{a}^2(M + \frac{m}{2}) - \bar{a}^2(M - \frac{m}{2})) \rho = 0 \quad (7)$$

We consider a chain of  $N$  molecules and impose cyclic boundary conditions. Then the Fourier transform of the two-point correlation function is defined by

$$F(k, M, t) = \sum_m e^{-ikm} \rho(M + \frac{m}{2}, M - \frac{m}{2}, t) \quad (8)$$

where  $k$  takes values in the first Brillouin zone (BZ),  $k \in (-\pi, \pi)$ . The inverse formula is <sup>1</sup>

$$\rho(M + \frac{m}{2}, M - \frac{m}{2}, t) = \frac{1}{M} \sum_k^{BZ} e^{ikm} F(k, M, t) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ikm} F(k, M, t) dk. \quad (9)$$

For  $m = 0$  one obtains

$$\bar{a}^2(N, t) = \frac{1}{N} \sum_k^{BZ} F(k, M, t) = \frac{1}{2\pi} \int_{-\pi}^{\pi} F(k, M, t) dk. \quad (10)$$

Now Fourier transforming equation (7) we get

$$\begin{aligned} & \frac{\partial F}{\partial t} + 4\lambda \sin k \sinh \frac{1}{2} \frac{\partial}{\partial M} F + \\ & 4\mu \sum_{j=1}^{\infty} \frac{(-1)^{j\pm 1}}{(2j-1)! 2^{2j-1}} \left( \frac{\partial^{2j-1}}{\partial M^{2j-1}} \bar{a}^2(M) \right) \left( \frac{\partial^{2j-1}}{\partial k^{2j-1}} F(k, M) \right) = 0 \end{aligned} \quad (11)$$

which is the expected nonlinear evolution equation for  $F(k, M, t)$  in a mixed configuration-wave number space  $(M, k)$ . We remark that  $F(k, M, t)$  is a periodic function in the reciprocal space,  $F(k + 2\pi) = F(k)$ .

---

<sup>1</sup>In dealing with Fourier series we are using  $\frac{1}{N} \sum_{m=1}^N e^{i(k_1-k)m} = \delta_{k, k_1}$  and  $\frac{1}{N} \sum_k^{BZ} e^{ik(m_1-m)=\delta_{m, m_1}}$ . Also the sum over  $k \in 1BZ$  can be transformed into an integral using the relation  $\frac{1}{N} \sum_k^{BZ} \dots \rightarrow \frac{1}{2\pi} \int_{-\pi}^{\pi} \dots dk$ .

### 3 Stability analysis

As the unperturbed problem we shall consider a basic solution  $F_0(k)$  independent of  $M$  and  $t$ . This is the random counterpart of the Stokes wave in a deterministic approach. A small perturbation around this homogeneous background is considered, namely

$$F(k, M, t) = F_0(k) + \epsilon F_1(k, M, t) \quad (12)$$

According to (10) we have also

$$\bar{a}^2(M, t) = \bar{a}_0^2 + \epsilon \bar{a}_1^2(M, t) \quad (13)$$

where

$$\begin{aligned} \bar{a}_0^2 &= \frac{1}{N} \sum_k^{BZ} F_0(k) = \frac{1}{2\pi} \int_{-\pi}^{\pi} F_0(k) dk \\ \bar{a}_1^2(M, t) &= \frac{1}{N} \sum_k^{BZ} F_1(k, M, t) = \frac{1}{2\pi} \int_{-\pi}^{\pi} F_1(k, M, t) dk \end{aligned} \quad (14)$$

When (12) is introduced into (11), neglecting terms of order  $\epsilon^2$ , the following linear evolution equation for  $F_1$  is obtained

$$\begin{aligned} &\frac{\partial F_1}{\partial t} + 4\lambda \sin k \sinh \frac{1}{2} \frac{\partial}{\partial M} F + \\ &4\mu \sum_{j=1}^{\infty} \frac{(-1)^{j+1}}{(2j-1)! 2^{2j-1}} \frac{\partial^{2j-1} F_0}{\partial k^{2j-1}} \frac{\partial^{2j-1} \bar{a}_1^2(M)}{\partial M^{2j-1}} = 0 \end{aligned} \quad (15)$$

Looking for a plane wave solution

$$F_1(k, M, t) = f_1(k) e^{i(KM - \Omega t)}$$

after little algebra the following stability integral equation is found

$$1 + \frac{\mu}{4\pi\lambda \sin \frac{K}{2}} \int_{-\pi}^{\pi} \frac{F_0(k + \frac{K}{2}) - F_0(k - \frac{K}{2})}{\sin k - \frac{\Omega}{4\lambda \sin \frac{K}{2}}} dk = 0 \quad (16)$$

The modulational instability is related to  $\Omega$  complex with a positive imaginary part,  $Im\Omega > 0$ . It is convenient to compare (16) with the similar result for the continuum case of the nonlinear Schrödinger equation [8]

$$1 + \frac{\mu}{K\omega_2} \int_{-\infty}^{\infty} \frac{F_0(k + \frac{K}{2}) - F_0(k - \frac{K}{2})}{k - \frac{\Omega}{2K\omega_2}} dk = 0 \quad (17)$$

Although there are significant differences between the two expressions, when the width of the spectrum  $F_0(k)$  is vanishingly small, the final results will look very similar.

### 3a. Gaussian spectrum

As a first example let us assume  $F_0(k)$  to be a Gaussian function

$$F_0(k) = \frac{\sqrt{2\pi}}{\sigma} \bar{a}_0^2 e^{-\frac{k^2}{2\sigma^2}}. \quad (18)$$

This expression doesn't satisfy the periodicity condition  $F_0(k + 2\pi) = F_0(k)$ , but for  $\sigma$  vanishingly small the errors introduced are negligible. Also the relation (14) is satisfied up to exponentially small terms.

It is convenient to introduce the new integration variable  $t = \frac{1}{\sqrt{2}\sigma}(k \pm \frac{K}{2})$  and the notations

$$z_{\pm} = \frac{1}{\sqrt{2}\sigma} \left( \frac{\Omega}{2\lambda \sin K} \pm \tan \frac{K}{2} \right) \quad (19)$$

$$f_{\pm}(t) = \frac{1}{\sqrt{2}\sigma} \left( \sin \sqrt{2}\sigma t \pm \tan \frac{K}{2} (1 - \cos \sqrt{2}\sigma t) \right).$$

Then (16) becomes

$$\frac{\bar{a}_0^2}{\sqrt{2\pi}\sigma} \frac{\mu}{\lambda \sin K} \int_{-\frac{\pi}{\sqrt{2}\sigma}}^{\frac{\pi}{\sqrt{2}\sigma}} e^{-t^2} \left( \frac{1}{z_+ - f_+} - \frac{1}{z_- - f_-} \right) dt = 1. \quad (20)$$

When  $\sigma \ll 1$  the integration limits can be extended to infinity and  $f_{\pm}$  can be expanded in Taylor series as

$$f_{\pm} = t \pm \left( \frac{\sigma}{\sqrt{2}} \tan \frac{K}{2} \right) t^2 - \left( \frac{\sigma^2}{3} \right) t^3 + O(\sigma^3).$$

Also  $z_{\pm} \gg 1$  and the integral in (20) can be written as

$$\int_{-\infty}^{\infty} e^{-t^2} \left( \frac{1}{z_+} \left( 1 + \frac{f_+}{z_+} + \frac{f_+^2}{z_+^2} + \dots \right) - \frac{1}{z_-} \left( 1 + \frac{f_-}{z_-} + \frac{f_-^2}{z_-^2} + \dots \right) \right) dt$$

and integrated term by term. Neglecting terms of order  $\sigma^3$  we get

$$1 = \bar{a}_0^2 \frac{\mu}{\lambda \sin K} \left[ \frac{1}{\frac{\Omega}{2\lambda \sin K} + \tan \frac{K}{2}} \left( 1 + \frac{\sigma^2}{\frac{\Omega}{2\lambda \sin K} + \tan \frac{K}{2}} \left( \frac{1}{2} \tan \frac{K}{2} + \frac{1}{\frac{\Omega}{2\lambda \sin K} + \tan \frac{K}{2}} \right) \right) - \frac{1}{\frac{\Omega}{2\lambda \sin K} - \tan \frac{K}{2}} \left( 1 + \frac{\sigma^2}{\frac{\Omega}{2\lambda \sin K} - \tan \frac{K}{2}} \left( -\frac{1}{2} \tan \frac{K}{2} + \frac{1}{\frac{\Omega}{2\lambda \sin K} - \tan \frac{K}{2}} \right) \right) \right] \quad (21)$$

Retaining only the terms independent of  $\sigma$  and considering  $\Omega$  purely imaginary,  $\Omega = i\Omega_i$ , one obtains

$$\Omega_i = 4\lambda \sin \frac{K}{2} \sqrt{\bar{a}_0^2 \frac{\mu}{\lambda} - \sin^2 \frac{K}{2}} \quad (22)$$

Modulational instability corresponds to  $\Omega_i > 0$  and is obtained if  $\mu$  and  $\lambda > 0$  and if  $\sin^2 \frac{K}{2} < \bar{a}_0^2 \frac{\mu}{\lambda}$ .

### 3b. Lorentzian spectrum

A simpler example is a Lorentzian form for  $F_0(k)$

$$F_0(k) = \bar{a}_0^2 \frac{2p}{k^2 + p^2}. \quad (23)$$

One assumes  $p \ll 1$  and then (14) is approximately satisfied.

Let us introduce the new integration variable  $t = \frac{1}{p}(k \pm \frac{K}{2})$  and the notations

$$z_{\pm} = \frac{1}{p} \left( \frac{\Omega}{4\lambda \sin \frac{K}{2}} \pm \sin \frac{K}{2} \right) \quad (24)$$

$$f_{\pm} = \frac{1}{p} \left[ \cos \frac{K}{2} \sin pt \pm \sin \frac{K}{2} (1 - \cos pt) \right]$$

If  $p \ll 1$  the integration limits can be extended to infinity and (16) becomes

$$1 = \frac{\mu}{2\pi \lambda \sin \frac{K}{2}} \int_{-\infty}^{\infty} \frac{dt}{1+t^2} \left( \frac{1}{z_+ - f_+} - \frac{1}{z_- - f_-} \right) \frac{1}{p}. \quad (25)$$

The integral can be done in the complex  $t$ -plane, closing the contour in the lower half-plane where one pole exists at  $t = -i$ .<sup>2</sup> The result is

$$1 = \frac{\mu}{\lambda \cos^2 \frac{K}{2}} \bar{a}_0^2 \frac{\cosh p}{\left( \frac{\mu}{\lambda \cos^2 \frac{K}{2}} \bar{a}_0^2 - \tan^2 \frac{K}{2} \cosh p \right)^2 + \tan^2 \frac{K}{2} \cosh^2 p}$$

so

$$\Omega_i = 2\lambda \sin K \left( \sqrt{\frac{\mu}{\lambda \cos^2 \frac{K}{2}} \bar{a}_0^2 \cosh p - \tan^2 \frac{K}{2} \cosh^2 p} - \sinh p \right) \quad (26)$$

When only terms linear in  $p$  are considered

$$\Omega_i = 4\lambda \sin \frac{K}{2} \left( \sqrt{\frac{\mu}{\lambda} \bar{a}_0^2 - \sin^2 \frac{K}{2}} - p \cos \frac{K}{2} \right). \quad (27)$$

Modulational instability occurs if  $\sin \frac{K}{2} < \sqrt{\frac{\mu}{\lambda} \bar{a}_0^2}$  and if  $p \cos \frac{K}{2} < \sqrt{\frac{\mu}{\lambda} \bar{a}_0^2 - \sin^2 \frac{K}{2}}$ .

Both results (22) and (27) can now be compared with the similar results obtained for the NLS equation [8], namely

$$\begin{aligned} \Omega_i^{(G)} &= 2K\omega_2 \sqrt{\frac{\mu}{\omega_2} \bar{a}_0^2 - \frac{K^2}{4}} \\ \Omega_i^{(L)} &= 2K\omega_2 \left( \sqrt{\frac{\mu}{\omega_2} \bar{a}_0^2 - \frac{K^2}{4}} - p \right) \end{aligned} \quad (28)$$

when the superscripts  $G/L$  refers to a Gaussian/Lorentzian spectrum. It is easily seen that (28) are the long wave limit ( $K \ll 1$ ) of (22) and (27) respectively.

In the case of a Lorentzian spectrum, like in the case of NLS equation, if  $p$  in (27) is greater than the square root,  $\Omega_i$  becomes negative and no instability can develop. This situation is similar with the well known phenomenon of Landau damping in plasma physics [13], [14]. The relation (27) contains two competing processes. The first is the generation of sidebands around

---

<sup>2</sup>The integrand has a second pole, coming from the zeroes of  $\sin k - iz = 0$ , in the  $k$ -complex lower half-plane,  $k = \pm\pi - ik_i$ ,  $k_i = \arg \sinh z$ ,  $z = \frac{\Omega_i}{4\lambda \sin \frac{K}{2}}$ , but its contribution can be neglected in the first order.



the carrier wave and their resonant interaction, process responsible for the square root in (27) and for the MI of the initial constant amplitude wave solution, while the second is related to how short are the correlations in the initial condition. For a Lorentzian form of  $F_0(k)$  the unperturbed two-point correlation function in the initial stage is given by  $\rho_0(n_1, n_2) = \bar{a}_0^2 e^{-p|n_1 - n_2|}$  and if it is of too short range it opposes to the MI.

In conclusion a complete discrete discussion of the randomness effects on the MI of the self trapping equation was done and the discrete effects are easily seen in the final results, compared with the similar ones found for the NLS equation.

*Helpful discussions with Dr. A.S. Cârstea are kindly acknowledged. This research was supported under the contract 66, CERES Programme, with the Ministry of Education and Research.*

## References

- [1] J.C. Eilbeck, P.S. Lomdahl, A.C. Scott, *Physica D* **16**, 318 (1985)
- [2] *Davydov's Soliton Revisited. Self Trapping of Vibrational Energy in Proteins*, edited by P.L. Christiansen, A.C. Scott, NATO ASI Series B **243** (Plenum Press, New York, 1990)
- [3] A.S. Davydov, N.I. Kislukha, *phys.stat.sol.(b)* **59**, 465 (1973); **75**, 735 (1976);  
A.S. Davydov, *Physica Scripta* **20**, 387 (1979)
- [4] A.S. Davydov, *Solitons in Molecular Systems* (Reidel, Dordrecht, 1985)
- [5] L. Cruzeiro-Hansson, H. Feddersen, R. Flesch, P.L. Christiansen, M. Salerno, A.C. Scott, *Phys. Rev. B* **42**, 522 (1990)
- [6] T.B. Benjamin, J.E. Feir, *J. Fluid Mech.* **27**, 417 (1967)
- [7] R.K. Dodd, J.C. Eilbeck, J.D. Gibbon, H.C. Morris, "Solitons and Non-linear Wave Equations" (Acad. Press, 1982)

- [8] D. Grecu, Anca Vişinescu, "Modulational instability in some nonlinear one-dimensional lattices and soliton generation" (to be published in Ann.Univ. Craiova)
- [9] I.E. Alber, Proc. Roy. Soc. London A **363**, 525 (1978)
- [10] M. Onorato, A. Osborne, M. Serio, R. Fedele, nlin.CD/0202026;  
R. Fedele, P.K. Shukla, M. Onorato, D. Anderson, M. Lisak, nlin.CD/0207050
- [11] C.M. Gardiner "Handbook of Stochastic Methods" (Springer Verlag, Berlin, 1983)
- [12] E. Wigner, Phys. Rev. **40**, 749 (1932)  
J.E. Moyal, Proc. Cambridge Phyl. Soc. **45**, 99(1949)
- [13] A. Simon, in "Plasma Physics" p. 163 (IAEA, Vienna, 1965);  
L.D. Landau, J. Phys. USSR **10**, 25 (1946)
- [14] M.J. Ablowitz, H. Segur "Solitons and the Inverse Scattering Transform", see appendix (SIAM, Philadelphia, 1981)