

ИЗДАТЕЛЬСТВО  
МЕДИЦИНСКОЙ  
АКАДЕМИИ  
УБЕНА



75-4-179  
高工研図書室

S. Berceanu

E2 - 8542

ON TAUBERIAN THEOREMS  
AND INCLUSIVE PROCESSES

1975

Издательский отдел Объединенного института ядерных исследований  
Заказ 19236. Тираж 690. Уч.-изд. листов 0,67.  
Редактор Э.В. Пашкевич. Подписано к печати 5.3.75 г.

Берчану Ш.

E2 - 8542

Тауберовы теоремы и инклюзивные процессы

Показано, что дифракционные процессы не могут насыщать кинематическую границу средней множественности. Предполагая, что граница независима от энергии и существуют общие условия для поведения в центральной области, для структурной функции получаем ненасыщение границы  $\langle n \rangle$  и  $\langle p_{\perp} \rangle$ . Если  $\langle n \rangle$  и  $\langle p_{\perp} \rangle$  растут медленнее, чем  $\sqrt{s}$ , наблюдается отдельное сохранение продольного импульса для каждой полусферы.

Сообщение Объединенного института ядерных исследований  
Дубна 1975

Berceanu S.

E2 - 8542

On Tauberian Theorems and Inclusive Processes

It is argued that diffractive processes cannot lead to the saturation of the kinematical bound of the average multiplicity. If the structure functions for inclusive reactions have scaling upper bounds and verify some general conditions at large transverse momentum in the central region, then the unsaturation of the kinematical bound of  $\langle p_{\perp} \rangle$  and  $\langle n \rangle$  occurs. It is shown also that if  $\langle p_{\perp} \rangle$  and  $\langle n \rangle$  increase asymptotically slower than  $\sqrt{s}$ , then a left-right conservation of reduced c.m. longitudinal momentum follows.

Communication of the Joint Institute for Nuclear Research  
Dubna 1975

### RANGE OF JINR PUBLICATIONS

The preprints and communications of the Joint Institute for Nuclear Research (JINR) are considered to be original publications. They are published in accordance with Article 4 of the JINR Statut. Difference between the preprints and communications consists in that text of the preprint will be published in future in one of scientific journals or in aperiodic collections.

### Indexing

The preprints, communications and deposited publications of the JINR have a single numbering (four last figures of the index).

The first sign of the index - a letter, denotes the language the paper is published in:

"P" - published in Russian;

"E" - published in English;

"D" - published both in Russian and English.

The preprints and communications, which are distributed only to the Member States of the JINR, have no letter-index.

The figure following the letter denotes the subject category of the given publication. List of subject categories of the JINR publications is sent periodically to the recipients.

Above-described index is placed in the right upper corner of the cover and on the title-page of each publication.

### References

In bibliographical references to the JINR preprints and communications we recommend to indicate: author's initials and name, an abbreviation of the name of the institute, index, place and year of publication.

An example of bibliographical reference:

I. I. Ivanov. JINR, P2-4985, Dubna, 1971.

S. Berceanu\*

## ON TAUBERIAN THEOREMS AND INCLUSIVE PROCESSES

\*Present address: Institute of Atomic Physics, Bucharest.

The experimental data on multiparticle production at high energy indicate that the values of the total average multiplicity and average transverse momentum are systematically smaller than the phase space allows<sup>1/</sup>. However, theoretically it is not clear if the general principles (like unitarity and analyticity) control the behaviour of the average multiplicity<sup>2,3</sup> $\langle n \rangle$  and average transverse momentum  $\langle p_{\perp} \rangle$ . Therefore, it is useful to find general conditions for the unsaturation of kinematical limits of  $\langle n \rangle$  and  $\langle p_{\perp} \rangle$ .

In this note, firstly we show that the diffractive processes cannot lead to the saturation of the kinematical bound of  $\langle n \rangle$  (Sect. 1). In Sect. 2 we argue that scaling<sup>4/</sup> upper bounds for the structure function of the inclusive reactions implies the unsaturation of the kinematical bound of  $\langle n \rangle$  and  $\langle p_{\perp} \rangle$ . Further, if  $\langle n \rangle$  and  $\langle p_{\perp} \rangle$  increase asymptotically slower than  $\sqrt{s}$ , then a separate conservation of the reduced center of mass (c.m.) energy and longitudinal momentum occurs on both c.m. hemispheres, where  $\sqrt{s}$  denotes the c.m. energy. The results are also extended to  $m$ -particle inclusive reactions (Sect. 3).

The proofs in this paper are mainly based on applications of some Tauberian theorems which seem to be a powerful tool for dealing with scaling properties.

### 1. Average Multiplicity of Diffractive Processes

We shall prove that the diffractive processes do not lead to the saturation of kinematical bound of  $\langle n \rangle$ .

Let us denote by  $\sigma_n$  the (exclusive)  $n$ -particle production cross section and by  $\sigma_{tot}$  the total cross section of

a hadronic ab collision. Introducing the probabilities for a -particle production

$$P_n(s) = \sigma_n(s) / \sigma_{tot}(s),$$

the total average multiplicity is defined as

$$\langle n(s) \rangle = \sum_{n=2}^{\infty} n P_n(s). \quad (1)$$

In fact, in the sum from the r. h. s. of eq. (1)  $n$  runs till the integer part of  $(\sqrt{s} - m_a - m_b) / m_\pi$ , but we express  $\sqrt{s}$  in pionic masses and neglect the masses  $m_a, m_b$  of the colliding particles  $a, b$ , respectively.

By diffractive processes we mean processes where

$$\lim_{s \rightarrow \infty} P_n(s) = p_n > 0. \quad (2)$$

The average multiplicity of a diffractive process is obviously  $\sqrt{s}$

$$\langle n_D \rangle = \sum_{n=2}^{\infty} n P_n, \quad (3)$$

where  $P_n$  are defined by eq. (2).

Now we shall use a Tauberian theorem (due to Ikebara/5/). If the Dirichlet series

$$D(\xi) = \sum_{n=2}^{\infty} a_n / n^\xi, \quad a_n > 0 \quad (4)$$

is convergent for  $\text{Re } \xi > 1$  and regular at  $\xi = 1$ , then

$$\lim_{x \rightarrow \infty} x^{-1} \sum_{n < x} a_n = 0. \quad (5)$$

Taking in the Dirichlet series (4)  $a_n = n P_n$ , the conditions of the Ikeara theorem are fulfilled because

$$D(1) = \sum_{n=2}^{\infty} P_n \leq 1.$$

Hence, with  $x = \sqrt{s}$  in eq. (5) we obtain

$$\lim_{s \rightarrow \infty} \sum_{n=2}^{\sqrt{s}} n P_n / \sqrt{s} = 0,$$

i.e.,

$$\lim_{s \rightarrow \infty} \langle n_D \rangle / \sqrt{s} = 0, \quad (6)$$

and so we have proved our assertion. We remember that some models<sup>3/</sup> claim the saturation of the kinematical bound of  $\langle n \rangle$ . Eq. (6) shows that if the saturation occurs, then it could come only from non-diffractive processes. Details will be discussed elsewhere<sup>6/</sup>. Here we want to stress also the utility of another Tauberian theorem<sup>7/</sup> for diffractive processes. This theorem enables to establish (rigorously) the expected asymptotic behaviour of the multiplicity moments,

$$\langle n^k \rangle \sim s^{(k-1)/2}, \quad k = 2, 3, \dots, \quad (7)$$

for diffractive models<sup>8/</sup> for which

$$P_n \sim 1/n^2. \quad (8)$$

We express the Tauberian theorem<sup>7/</sup> in a sufficient form for the present purpose. Let us suppose that the series

$$Q(t) = \sum_{m=0}^{\infty} q_m t^m, \quad q_m \geq 0 \quad (9)$$

converges for  $0 \leq t < 1$ . If  $L$  varies slowly<sup>7/</sup> at  $\infty$ ,  $0 \leq \rho < \infty$ , and the sequence  $\{q_m\}$  is monotonic, then the relation

$$\sum_{m=0}^N q_m \sim \frac{1}{\Gamma(\rho+1)} N^\rho L(N), \quad N \rightarrow \infty \quad (10)$$

is equivalent to

$$q_N \sim \frac{1}{\Gamma(\rho)} N^{\rho-1} L(N), \quad N \rightarrow \infty. \quad (11)$$

The series (9) with

$$q_m = m^k P_m, \quad k = 2, 3, \dots \quad (12)$$

is evidently convergent for  $0 \leq t < 1$ . We proceed to evaluate the multiplicity moments for diffractive processes

$$\langle n^k \rangle = \sum_{m=2}^{\sqrt{s}} q_m = \sum_{m=2}^{\sqrt{s}} \sum_{k=2,3,\dots}^m p_m, \quad (13)$$

Supposing for  $p_m$  the behaviour (8),  $q_m$  (eq.(12)) takes the form (11) for  $\rho=k-1(N=\sqrt{s})$ . Then eq. (10) furnishes the behaviour (7) for the multiplicity moments (13) of diffractive processes.

## 2. Unsaturation of the Kinematical Bound of $\langle n \rangle$ and $\langle p_{\perp} \rangle$

We begin with some notations and definitions. Let us denote by  $f_c(x, p_{\perp}^2, s)$ ,

$$f_c(x, p_{\perp}^2, s) = \frac{2\pi}{\sigma_{tot}(s)} x_c^0 \frac{d\sigma_c}{dx dp_{\perp}^2}, \quad (14)$$

the structure function of the inclusive reaction

$$a + b \rightarrow c + \text{anything}. \quad (15)$$

Here  $x$  is the reduced c.m. longitudinal momentum ( $p_{\parallel}$ ),  $p_{\perp}$  is the transverse momentum and  $x^0$  the reduced c.m. energy  $E$  of the particle  $c$  with mass  $M_c$ ,

$$x_c^0 = 2E_c/\sqrt{s} = 2\sqrt{p_{\parallel}^2 + p_{\perp}^2 + M_c^2}/\sqrt{s} = \sqrt{x^2 + 4(p_{\perp}^2 + M_c^2)}/s. \quad (16)$$

With these notations the total average multiplicity and total average transverse momentum can be written, respectively,

$$\langle n \rangle_1 = \sum_c \langle n_c \rangle = \sum_c \int \frac{f_c(x, p_{\perp}^2, s)}{x_c^0} dx dp_{\perp}^2, \quad (17)$$

$$\langle p_{\perp} \rangle_1 = \sum_c \langle p_{\perp} \rangle_c \langle n_c \rangle = \sum_c \int \frac{f_c(x, p_{\perp}^2, s)}{x_c^0} p_{\perp} dx dp_{\perp}^2. \quad (18)$$

The index 1 specifies that we deal with average values

for the one-particle inclusive reaction (15). Analogously are introduced the total average of  $p_{\perp}^k$ ,  $k=2,3,\dots$ . The generalization to  $m$ -particle inclusive reactions is sketched in Sect. 3.

A useful quantity in the following is the average

$$\langle p_{\perp}^k x^0 \rangle_1 = \sum_c \int p_{\perp}^k f_c(x, p_{\perp}^2, s) dx dp_{\perp}^2, \quad k=0,1,2,\dots, \quad (19)$$

We need also the energy and longitudinal momentum sum rules<sup>9)</sup>, which in the present notations take the form

$$\langle x^0 \rangle_1 = 2, \quad (20)$$

$$\langle x \rangle_1 = 0. \quad (21)$$

With the notations

$$x_{\perp} = 2p_{\perp}/\sqrt{s}, \quad (22)$$

$$\langle N^k \rangle_1 = \langle n^k \rangle_1 / \sqrt{s}, \quad k=1,2,\dots, \quad (23)$$

our purpose is to find conditions in which the following relations hold:

$$\lim_{s \rightarrow \infty} \langle N \rangle_1 = 0, \quad (24)$$

$$\lim_{s \rightarrow \infty} \langle N_{\perp} \rangle_1 = 0. \quad (25)$$

Now let us suppose that the structure function (14) has the scaling bound

$$\int f_c(x, p_{\perp}^2, s) dx \leq g_c(p_{\perp})/2p, \quad (26)$$

and in the central region has the behaviour

$$\max_{x \in [-\epsilon, \epsilon]} f_c(x, p_{\perp}^2, s) \leq \frac{\gamma}{(p_{\perp}^2 + M_c^2)^{\beta} [\beta n(p_{\perp}^2 + M_c^2)]^{\beta}}, \quad \beta = 2 + \delta, \delta > 0. \quad (27)$$

It can be noted that the upper bound (27) for the behaviour at large  $p_{\perp}$  in the central region is a weak condition

for the structure function, which is verified by usual models/<sup>10</sup> for large  $p_{\perp}$ .

We assert that if the structure functions (14) have the scaling upper bounds (26) and verify the conditions (27) at large  $p_{\perp}$ , then the total average multiplicity and total average transverse momentum increase slower than  $\sqrt{s}$  when  $s \rightarrow \infty$  (i.e., the relations (24), (25) hold). Also, if the condition (26) is verified, then

$$\lim_{x \rightarrow \infty} \langle x_{\perp}^2 \rangle = 0. \tag{28}$$

For the proof the following Tauberian theorem/<sup>11</sup> will be used. If

$$\lim_{x \rightarrow \infty} \int_1^x \frac{b(y)}{y} \left(1 - \frac{y}{x}\right) dy = a, \tag{29}$$

where  $b$  is bounded real function, then

$$\lim_{x \rightarrow \infty} \int_1^x \frac{b(y)}{y} dy = a. \tag{30}$$

Introducing in eq. (20) the bound (26) and applying the Tauberian theorem (29)-(30) with  $b(y) = y g_c(y)$  and  $x = \sqrt{s}$ , we get readily

$$\lim_{s \rightarrow \infty} \int_0^{\sqrt{s}/2} dp_{\perp} p_{\perp} g_c(p_{\perp}) / \sqrt{s} = 0. \tag{31}$$

With the definitions (16) and (19) for  $k = 1$ , and the bound (26), eq. (28) is obtained immediately. Further, with the bounds (27) for  $f_c(x, p_{\perp}^2, s)$  in the central region and eq. (31) in the fragmentation region, it can be verified that the relations (24) and (25) hold.

### 3. Asymptotic Left-Right Conservation of $x$ and $x^0$

In this Section it will be supposed that eqs. (24) and (25) hold. We argue that separate left-right conservation of the reduced c.m. energy and longitudinal momentum occurs. Also we shall be concerned with the evaluation

of the asymptotic behaviour of more general average quantities for the  $m$ -particle inclusive reactions

$$a + b \rightarrow c_1 + \dots + c_m + \text{anything}, \tag{32}$$

Let us consider the average of a function  $F_{c_1 \dots c_m}(\vec{p}_1, \dots, \vec{p}_m)$  on a given volume of the phase space

$$\langle F(R) \rangle_m = \sigma_{\text{tot}}^{-1}(s) \sum_{c_1, \dots, c_m} \int_R F_{c_1 \dots c_m}(\vec{p}_1, \dots, \vec{p}_m) \frac{d\sigma_{c_1 \dots c_m}}{dp_1 \dots dp_m} \tag{33}$$

where  $\vec{p}_i$  denotes the 3-momentum vector of the particle  $c_i$  ( $i = 1, \dots, m$ ) and  $d\sigma_{c_1 \dots c_m} / dp_1 \dots dp_m$  the differential cross section of the process (32). In the following we take  $F$  as the monoms

$$F_{c_1 \dots c_m} = \prod_{i=1}^m (p_{\perp i})^{\alpha_i} (p_{\perp i})^{\beta_i} (E_i)^{\gamma_i}, \alpha_i, \beta_i, \gamma_i > 0. \tag{34}$$

If in eq. (33) we introduce  $F$  with particular values  $\alpha, \beta, \gamma$  from eq. (34), we get the averages (17)-(21). When  $F_{c_1 \dots c_m} = 1$  ( $m > 1$ ), we get the binomial multiplicity moments in the  $R$  region of the phase space

$$\langle \nu(R) \rangle_m = \sigma_{\text{tot}}^{-1}(s) \sum_{c_1, \dots, c_m} \int_R \frac{d\sigma_{c_1 \dots c_m}}{dp_1 \dots dp_m} \tag{35}$$

It may be remarked that the inequality

$$|p_{\perp}| \leq E_c \leq |p_{\perp}| + p_{\perp} + M_c \tag{36}$$

implies the same asymptotic behaviour of  $x^0$  and  $x$  in the average (33) of the monoms (34), when eq. (25) holds.

Further we list 4 consequences of eqs. (24), (25) for various averages (33)-(34).

- a. If eqs. (24), (25) hold, then

$$\lim_{s \rightarrow \infty} \langle x(r/l) \rangle_1 = 1, \quad (37)$$

i.e., if the total average multiplicity and total average transverse momentum increase slower than  $\sqrt{s}$  when  $s \rightarrow \infty$ , then a separate right (left) conservation of the reduced c.m. longitudinal momentum (and energy) occurs.

Adding the sum rules (20) and (21) and taking into account ineq. (36), we get easily

$$\langle x(r) \rangle_1 + \langle x \rangle_1 > 1 + \sum_c M_c \langle n_c \rangle / \sqrt{s} \geq 1 \geq \langle x(r) \rangle_1.$$

If eqs. (24), (25) are true, then eq. (37) follows.

b. If eq. (25) holds, then

$$\lim_{s \rightarrow \infty} \langle x \rangle_1 = \begin{cases} \beta, & \text{when } \beta > 0 \\ \lambda, & \text{when } \beta = 0, \end{cases} \quad (38)$$

where

$$\lambda = \begin{cases} \gamma, & \text{if } \gamma = 0, 1 \\ 0 < \lambda < 1, & \text{if } 0 < \gamma < 1. \end{cases} \quad (39)$$

Here by  $\gamma$  we denote

$$\gamma = \lim_{s \rightarrow \infty} \langle x^2 \rangle_1.$$

The behaviour from eqs. (38), (39) can be checked applying the Schwartz inequality.

c. If eq. (24) holds, then

$$\lim_{s \rightarrow \infty} \langle N^k \rangle_1 = 0, \quad k = 1, 2, \dots; \quad \lim_{s \rightarrow \infty} \langle r \rangle_1 / s^{m/2} = 0, \quad (40)$$

where  $N$  is defined by eq. (23) and  $r$  by eq. (35).

d. If eqs. (24), (25) hold, then

$$\lim_{s \rightarrow \infty} \frac{\langle p_{\perp,1}^k \rangle_1}{(\sqrt{s}/2)^{m+k}} = 0, \quad m > 1, \quad k \geq 1, \quad (41)$$

where the index 0 refers to the energy.

For proving eq. (41), the energy-momentum sum rules<sup>9</sup> for  $m$ -particle inclusive reactions are needed:

$$(P - p_{c_1} - \dots - p_{c_m}) \frac{d\sigma_{c_1 \dots c_m}}{dp_1 \dots dp_{m-1}} = \sum_c \int p_{c_m} \frac{d\sigma_{c_1 \dots c_m}}{dp_1 \dots dp_{m-1}} dp_1 \dots dp_m,$$

where  $p_{c_i}$  denotes the energy-momentum 4-vector of the particle  $c_i$  ( $i=1, \dots, m$ ) and  $P$  is the energy-momentum 4-vector of the initial state.

In conclusion, the properties (37)-(41) are consequences of the relations (24) and (25). Using Tauberian theorems we succeeded in Sect. 2 to prove that the scaling properties (26) and (27) of the structure functions are sufficient conditions for eqs. (24), (25). A more general justification of eqs. (24) and (25) will be presented elsewhere<sup>6</sup>.

The author acknowledges Dr. A.C.Gheorghe for initiating him in the use of Tauberian theorems.

## References

1. D.R.O.Morrison. Proceedings of the Fourth International Conference on High-Energy Collisions, Oxford, (1972); D.R.O.Morrison. Review lectures given at the Fifth Hawaii Topical Conference in Particle Physics, CERN/D.Ph.II/PHYS 73-46, CERN preprint (1973).
2. N.N.Khuri. Phys.Rev., D9, 1802 (1974).
3. S.Berceanu, to be published.
4. R.P.Feynman. Phys.Rev.Lett., 23, 1414 (1969).
5. S.Leng. Algebraic numbers, Izdat. Mir, Moscow, §VIII (1966) (in Russian).
6. S.Berceanu, A.C.Gheorghe, to be published.
7. W.Feller, An introduction to probability theory and its applications, Izdat. Mir, Moscow, Vol. II, §XIII, 513 (1967) (in Russian).
8. C.Quigg, J.H.Wang, C.N.Yang. Phys.Rev.Lett., 28, 1290 (1972).
9. L.S.Brown. Phys.Rev., D5, 748 (1972).
10. S.D.Ellis, R.Thun. Ref. TH 1874-CERN, CERN preprint (1974).



11. N. Dunford, J. T. Schwartz. Linear operators, Part II, Izdat. Mir, Moscow, 165 (1966) (In Russian).

Received by Publishing Department  
on January 20, 1975.

### Conditions of Exchange

The preprints and communications of the Joint Institute for Nuclear Research are distributed free of charge on the mutual exchange basis to the universities, institutes, libraries, scientific groups and individual scientists of more than 50 countries.

We expect that the recipients of the JINR publications will have a possibility to display initiative in sending free of charge publications to Dubna. We receive, on the exchange basis, scientific books, journals, preprints and other types of publications on the subject categories of our Institute.

The only kind of publications that we do not require are the reprints of the articles already published in scientific journals.

In a number of cases we address to scientific institutions - the most prominent recipients of our publications - with the request to send us free of charge some books or subscribe for our library to the scientific journals published in their countries.

### Requests

The Publishing Department fulfills annually about 3000 individual requests for our preprints and communications. The index of our publication must be obligatory indicated in such requests.

### Addresses

Letters on all the questions concerning the exchange of publications as well as requests for individual publications are to be sent at the address:

Publishing Department  
Joint Institute for  
Nuclear Research  
Head Post Office,  
P.O. Box 79  
101000 Moscow,  
U.S.S.R.

We kindly ask to send all the publications on the exchange basis and also free of charge subscriptions to scientific journals at the address:

Scientific-Technical Library  
Joint Institute for  
Nuclear Research  
Head Post Office  
P.O. Box 79  
101000 Moscow,  
U.S.S.R.